

Journal of Physical Chemistry and Functional Materials

Home Page of Journal: <https://dergipark.org.tr/jphcfum>



Structure Determination of Serotonin using Quantum Approximations

Ibrahim Isah Nasidi^{a,*}, Mansur said^c and Yau Datti^b

^aDepartment of Physics, Faculty of Science, Firat University, 23169 Elazig Turkey

^bDepartment of Chemistry, Yusuf Maitama Sule University, Kano. Nigeria

^cDepartment of Physics, Yusuf maitama Sule, Kano. Nigeria

* Corresponding author: E-mail: sinonasidi@yahoo.com

ABSTRACT

The role played by computational quantum theory methods especially Hartree-Fock (HF) and Density functional theory (DFT) in physical chemistry and theoretical physics is very important and helpful. Serotonin is neurotransmitter and has medicinal value, that is why it is important to understand its stability, reactivity and structure analysis. Serotonin is a monoamine neurotransmitter. Its biological function is complex and multifaceted, modulating mood, cognition, reward, learning, memory, and numerous physiological processes such as vomiting. In this study, we optimize the energy band gaps using different basis sets for both DFT and HF methods. LanL2MB at DFT method was selected for the molecule. IR, NMR and UV were also investigated during the research.

ARTICLE INFO

Keywords:

Serotonin.

Band gap energy.

DFT.

Hartree-Fock Theory.

Received: 3-November-2020,

Accepted: 20-November-2020

ISSN: 2651-3080

1. INTRODUCTION

Approximately 90% of the human body's total serotonin is located in the enterochromaffin cells in the GI tract, where it regulates intestinal movements.[12][13] About 8% is found in platelets and 1%-2% in the CNS.[14] The serotonin is secreted luminally and basolaterally, which leads to increased serotonin uptake by circulating platelets and activation after stimulation, which gives increased stimulation of myenteric neurons and gastrointestinal motility.[15] The remainder is synthesized in serotonergic neurons of the CNS, where it has various functions. These include the regulation of mood, appetite, and sleep. Serotonin also has some cognitive functions, including memory and learning

Many researchers have focused on the development of a sensitive method for the measurement of serotonin which is important for certain diseases [12-14]. The resulting of nuclear magnetic resonance (NMR) and DFT was provided information in the understanding its neurotransmitter activity in aqueous solutions. DFT theoretical calculation with the basis set of LanL2MB was

analyzed various hydrogen bond of serotonin. In this study, we will explain and find the energy band gaps by using Gaussian program software. The different basis sets (3-21G, 6-31G, SDD, 6-311G, LanL2DZ, LanL2MB) for two methods (DFT and HF) performed on serotonin geometry. LanL2MB was chosen at the DFT method to show the IR and NMR spectra for the geometry structure.

2. COMPUTATIONAL METHODS

The serotonin structure was drawn by Chem. Bio Draw Ultra 14.0 (Fig. 1). The structure of structure was optimized by both density functional theory (DFT) and Hartree-Fock (HF) [9]. Different basis sets have been applied to find the energy band gaps of serotonin. The reason why different basis sets was used is that each basis sets involves different functionalities and characteristics. Table 1 shows the energy band gaps for the two methods of different basis sets. The geometry system of serotonin at the B3LYP (DFT), LanL2MB basis set has been chosen [9, 18]. The frequency and NMR were calculated to this basis set [19] The Gaussian 09 program has been used for

all the calculations. Also, the origin program was performed for drawing the IR as in figure 3.




Figure 1. Structure of Serotonin

3. RESULT AND DISCUSSION

3.1. Energy bandgap

The first task of the computational work is to detect the optimized molecular structure using Gaussian 09 program. The energy band gaps associated with basis sets (3-21G, 6-31G, SDD, 6-311G, LanL2DZ, LanL2MB) were listed in Table 1. The energy band gaps for the Hartree-Fock (HF) method have higher values compared with the density functional theory (DFT). The optimized structural parameter of serotonin with the LanL2MB was selected by the DFT method [9].

Table 1. The energy bandgap for different basis sets for both HF and DFT methods.

Basis sets	HF method Energy gaps (ev)	DFT method Energy gaps (ev)
3-21G	15.8642158	10.2319514
6-31G	15.9734933	10.42233058
6-311G	16.2579401	10.9985208
SDD	16.4171111	11.2536823
LanL2DZ	16.4049088	10.9879455
LanL2MB	11.6954023	5.42916552

3.2. Molecular Structure:

The molecular structure and numbering of atom in the serotonin molecule was shown in Figure 1. The 72 vibration modes are present in the molecule. The first task of computational work was optimization of the molecule. The Gauss programs viewed the number of atoms scheme. To calculate the geometric structure DFT was used.

3.3 Vibrational Assignments:

The vibrational frequency was calculated by the DFT method at LanL2MB basis sets, FT-IR Figure 2 was observed with various vibration have been present in Table 2. The fundamental functional group present in the molecule vibrated at different frequencies as discussed below.

3.2.1. C-C /C=C vibrations:

The vibration carbon-carbon stretching in a ring usually occurs in a region 1400-1600 cm⁻¹ [21]. In aromatic benzene ring the vibration carbon-carbon was occurring in the region 1420-1625 cm⁻¹. In the aromatic like benzene has six-member rings, there are two or more vibration are occur in a region due to the vibration of the skeletal, the strong vibration was started at 1500 cm⁻¹. Moreover, where the ring is conjugated by the groups the further vibration observed at 1580 cm⁻¹. In the present work, the carbon-carbon stretching vibration occurred in a region 1768.02-1839.39 cm⁻¹. The strong starching vibration of C=C in a ring occurred in a region 1579.42-1675.25 cm⁻¹. The vibration carbon-carbon single bond occurred in 1417.40-1488.18 cm⁻¹ it is symmetrical vibration, but the stronger vibration for C-C in benzene ring was observed in region 1488.18 cm⁻¹. In 595.44, 649.6, 723.66 and 731.78 cm⁻¹ C-C and C=C in a plane (in a ring) was rocking but in a region 884.88, 896.45 and 964.37 cm⁻¹ was symmetrical Stretching.

3.2.2. C-H vibrations:

Commonly aromatic compounds in a region 3473-3468 cm⁻¹ were observed multiple weak bonds due to the C-H stretching vibration of the aromatic rings [14]. The C-H bending vibration in a plan was observed in a region 990-1390 cm⁻¹ and the peak was weak intensity [5]. This bond of C-H in a plane was very useful to found the characterization of compounds [26]. When the carbon and hydrogen are in a plane interaction was vibration above 1200 cm⁻¹ usually moves in the opposite direction. [7, 8]. The C-H out of the plane usually occurs in a region 700-1000 cm⁻¹ [19, 10]. In our anthracene molecule the C-H symmetrical stretching vibration out of the plane was observed between the regions 3027.75-3094.4 cm⁻¹, but the C-H in a plane was symmetrically vibrated between the regions 3181.27-3236.82 cm⁻¹ this is an agreement with literature survey. In a region 3236.82- 526.53 cm⁻¹ all C-H out of the plan was rocking vibrations. Also, al hydrogen was rocking vibration in serotonin molecules in a region between 1181.68- 1256.21cm⁻¹.

3.2.3. C-N vibration

In this study of serotonin structure was shown the vibration of C-N in a region 996.46 cm⁻¹ was symmetrical and in a region, 1152.66 cm⁻¹ was scissoring.

Table 2. Various vibration of atoms calculated by DFT (LanL2MB)

St. NO	vibration assignments	observed frequency (cm ⁻¹)	St. NO	vibration assignments	observed frequency (cm ⁻¹)
1	(C-H) out	43.77	32	(all H) ro	1181.68
2	(C-H) out	80.72	33	(all H) ro	1200.46
3	(C-H) out	128.26	34	(all H) ro	1227.25
4	(C-H) out	191.71	35	(all H) out,ro	1256.21
5	(C-H) out	249.98	36	(C1-O8)Sy,s	1281.71
6	(O7-H17)ro	273.63	37	(C4-C12)Sy,s	1326.77
7	(N14-H15,16)ro	283.72	38	(C12-C13)Sy,s	1358.67
8	(N14-H15,16)ro (C-O)ro	297.55	39	(C-C)in,Sy	1383.2
9	(C-H)out,ro	312.8	40	(all H)out, ro	1399.39
10	(C-H) out,ro	369.13	41	(C-C)in,Sy	1417.4
11	(C-H) out,ro	428.62	42	(C-C)in,Sy	1427.62
12	(C-H) out,ro	458.24	43	(C-C)in,Sy	1488.18
13	(C-H) out,ro	473.85	44	(C13-H21,22)sc	1527.34
14	(C-H) out,ro	490.4	45	(C12-H19,20)sc	1531.68
15	(C-H) out,ro	526.53	46	(C=C)in,Sy	1579.42
16	(N14-H15,16)ro	569.12	47	(C=C)in,Sy	1666.41
17	(C-C,C=C)in,ro	595.44	48	(C=C)in,Sy	1675.25
18	(C-C,C=C)in,ro	649.6	49	(N14-H15,16)sc	1709.1
19	(C-C,C=C)in,ro	723.66	50	(C12-H19,20)sy	3027.75
20	(C-C,C=C)in,ro	731.78	51	(C13-H21,22)sy	3043.08
21	(C-H) ro	794.48	52	(C12-H19,20)asy	3074.14
22	(C-H) ro	844.21	53	(C13-H21,22)asy	3094.4
23	(C-H) ro	857.95	54	(C5-H9)sy	3181.27
24	(C12-C13)sy	884.88	55	(C3-H10)sy	3199.5
25	(C12-C13)sy	896.45	56	(C2-H11)sy	3236.82
26	(C12-C13)sy	964.37	57	(N14-H15,16)sy	3551.73
27	(C-H)in,sc	976	58	(O8-H18)sy	3626.92
28	(C12-H19,20)ro	1020.82	59	(N14- H15,16)asy	3672.03
29	(C13-N14)sy	1103.82	60		
30	(O7-H17)ro	1126.49			
31	(C13-N14)sc				

Abbreviation: out: out of the plane, ro: rocking, sy: symmetrical, asy: anti-symmetrical, in: in the plan, sc: Scissoring.

3.2.4. NMR analysis

The ¹³C NMR chemical shifts used to identify chemical compounds [35]. GIAO (Gauge Including Atomic Orbital) is the faster procedure to calculation upon the basis set used. A GIAO method was preferable for many aspects state of the subject [19]. NMR was used to calculation chemical shift for the title compound best on TMS HF/LanL2MB(d) GIAO, TMS B3LYP/LanL2MB(2d, P)GIAO and oxygen best on the H₂O B3LYP/LanL2MB(2d, P) GIAO, H₂O HF/LanL2MB(d) GIAO. Table 3 showed the ¹³C NMR for all references. According to TMS HF/LanL2MB(d) reference, the carbon was ordered from C3>C1>C4>C6>C5>C2>C12>C13 that means carbon number C3 was higher ppm lower filed but C13 was lower ppm higher filed. However, for TMS b3LYP/LanL2MB(2d, P) reference to carbon atoms was

HF/LanL2MB(d) GIAO for carbon atoms. Hydrogen best on TMS HF/LanLMB(d) GIAO, TMS B3LYP/LanLMB(2d, P)GIAO and oxygen best on the H₂O B3LYP/LanL2MB(2d, P) GIAO, H₂O HF/LanL2MB(d) GIAO. Table 3 showed the ¹³C NMR for all references. According to TMS HF/LanL2MB(d) reference, the carbon was ordered from C3>C1>C4>C6>C5>C2>C12>C13 that means carbon number C3 was higher ppm lower filed but C13 was lower ppm higher filed. However, for TMS b3LYP/LanL2MB(2d, P) reference to carbon atoms was

order from C6>C1>C3>C4 >C2>C5>C13>12 in this reference C6 was observed at higher ppm lower filed but C12 was lower ppm higher filed. For the reference, CH4 HF/6-31 G(d), C1>C6>C4>C3>C2>C5>C13>C12, carbon number one (C1) was found in higher ppm and carbon number twelve found in lower ppm higher filed.




Figure 2. FT-IR Spectrum of Serotonin

For H-NMR Table 4 used two references TMS HF/LanL2MB G(d) GIAO and TMS B3LYP/LanL2MB+G(2d, P) GIAO. According to the reference TMS HF/LanLMB (d) GIAO the observation hydrogen peak started from higher filed (H15) to the end of lower filed (H11).H11>H10>H9>H18>H22>H17>H21>H19>H20>H16>H15. Besides for reference TMS B3LYP/LanL2MBG(2d, P) GIAO the order of hydrogen peak is the same as the previse reference with a little be different in a lower ppm for all hydrogen atoms. The NMR for nitrogen shown in Table 5, the peak was observed at 19.98 ppm for TMS HF/6-31 G(d)GIAO, but for TMS B3LYP/LanL2MB(2d, P) GIAO was observed at 17.5ppm.

Table 3. The observed C NMR chemical shifts in ppm for serotonin

C. NO	NONE ppm	TMS HF/LanL2 MB(d) GIAO Ppm	TMS B3LYP/LanL2 MB(2d,P) GIAO ppm	CH4 HF/LanL2M B(d) GIAO ppm
C1	54	146	151.9	145
C2	80.9	120.7	102	118.3
C3	75	155	132.7	124.1
C4	55.7	145.8	123	133.9
C5	84	124	98.3	115
C6	56	144	153.9	143.1
C12	151	71	31.9	48.1
C13	149	69	33.8	50.1

Table 4. The observed H NMR chemical shifts in ppm for serotonin

H. NO	NONE ppm	TMS HF/LanL2MB(d) GIAO	TMS B3LYP/LanL2MB(2d,P) GIAO
-------	----------	---------------------------	---------------------------------

		ppm	ppm
H9	26.79	5.81	5.1
H10	26.1	6.5	5.78
H11	25.83	6.7	6.0
H15	33.2	- 0.6	- 1.31
H16	32.41	0.9	- 0.58
H17	29.85	2.75	2.05
H18	27.6	5.0	4.28
H19	30.25	2.31	1.61
H20	30.8	1.8	1.1
H21	30.05	2.55	1.58
H22	29.78	2.82	2.13

4. CONCLUSION

In this work, all basis sets with both HF and DTF methods investigated to calculation bandgap energy for serotonin molecule. Proposed the better basis set is LanL2MB according to the literature previous. The IR shows the vibration for all atoms that is good agreement with the kinds of literature. NMR has determined the identity of the molecular structure. The peaks of atoms in NMR were the relation with the types of references which was used on it. For example, C1 has different shifted in a ppm for all three references. The reference H2O HF/LanL2MB(d)GIAO and H2O B3LYP/LanL2MB(2d, P)GIAO show the oxygen's ppm and it is closed to each other.

REFERENCES

- [1] Calculated using Advanced Chemistry Development (ACD/Labs) Software V11.02 (©1994–2011 ACD/Labs)
- [2] Mazák K, Dóczy V, Kőkösi J, Noszál B (April 2009). "Proton speciation and microspeciation of serotonin and 5-hydroxytryptophan". *Chemistry & Biodiversity*. **6** (4): 578–90. doi:10.1002/cbdv.200800087. PMID 19353542. S2CID 20543931.
- [3] Pietra S (1958). "[Indolic derivatives. II. A new way to synthesize serotonin]". *Il Farmaco; Edizione Scientifica (in Italian)*. **13** (1): 75–9. PMID 13524273.
- [4] Erspamer V (1952). "Ricerche preliminari sulle indolalchilamine e sulle fenilalchilamine degli estratti di pelle di Anfibio". *Ricerca Scientifica*. **22**: 694–702.
- [5] Tammisto T (1967). "Increased toxicity of 5-hydroxytryptamine by ethanol in rats and mice". *Annales Medicinae Experimentalis et Biologiae Fenniae*. **46** (3, Pt. 2): 382–4. PMID 5734241.
- [6] Jones D (2003) [1917], Roach P, Hartmann J, Setter J (eds.), *English Pronouncing Dictionary*, Cambridge: Cambridge University Press, ISBN 978-3-12-539683-8
- [7] "Serotonin". *Dictionary.com Unabridged*. Random House.
- [8] "Serotonin". *Merriam-Webster Dictionary*.
- [9] Young SN (November 2007). "How to increase serotonin in the human brain without drugs". *Journal of Psychiatry & Neuroscience*. **32** (6): 394–9. PMC 2077351. PMID 18043762.
- [10] González-Flores D, Velardo B, Garrido M, González-Gómez D, Lozano M, Ayuso MC, Barriga C, Paredes SD, Rodríguez AB (2011). "Ingestion of Japanese plums

- (*Prunus salicina Lindl. cv. Crimson Globe*) increases the urinary 6-sulfatoxymelatonin and total antioxidant capacity levels in young, middle-aged and elderly humans: Nutritional and functional characterization of their content". *Journal of Food and Nutrition Research*. **50** (4): 229–236.
- [11] Schlienger RG, Meier CR (2003). "Effect of selective serotonin reuptake inhibitors on platelet activation: can they prevent acute myocardial infarction?". *American Journal of Cardiovascular Drugs : Drugs, Devices, and Other Interventions*. **3** (3): 149–62. doi:10.2165/00129784-200303030-00001. PMID 14727927. S2CID 23986530.
- [12] King MW. "Serotonin". *The Medical Biochemistry Page*. Indiana University School of Medicine. Retrieved 1 December 2009.
- [13] Berger M, Gray JA, Roth BL (2009). "The expanded biology of serotonin". *Annual Review of Medicine*. **60**: 355–66. doi:10.1146/annurev.med.60.042307.110802. PMC 5864293. PMID 19630576.
- [14] Kling A (2013). 5-HT2A: a serotonin receptor with a possible role in joint diseases (PDF) (Thesis). Umeå Universitet. ISBN 978-91-7459-549-9.
- [15] Yano JM, Yu K, Donaldson GP, Shastri GG, Ann P, Ma L, Nagler CR, Ismagilov RF, Mazmanian SK, Hsiao EY (April 2015). "Indigenous bacteria from the gut microbiota regulate host serotonin biosynthesis". *Cell*. **161** (2): 264–76. doi:10.1016/j.cell.2015.02.047. PMC 4393509. PMID 25860609.
- [16] Vanhoutte PM (February 1987). "Serotonin and the vascular wall". *International Journal of Cardiology*. **14** (2): 189–203. doi:10.1016/0167-5273(87)90008-8. PMID 3818135.
- [17] Prah A, Purg M, Stare J, Vianello R, Mavri J (September 2020). "How Monoamine Oxidase A Decomposes Serotonin: An Empirical Valence Bond Simulation of the Reactive Step". *The Journal of Physical Chemistry. B*. **124** (38): 8259–8265. doi:10.1021/acs.jpcb.0c06502. PMC 7520887. PMID 32845149.
- [18] Huser A, Rohwedder A, Apostolopoulou AA, Widmann A, Pfitzenmaier JE, Maiolo EM, Selcho M, Pauls D, von Essen A, Gupta T, Sprecher SG, Birman S, Riemensperger T, Stocker RF, Thum AS (2012). "The serotonergic central nervous system of the *Drosophila* larva: anatomy and behavioral function". *PLOS ONE*. **7** (10): e47518. Bibcode:2012PLoS...747518H. doi:10.1371/journal.pone.0047518. PMC 3474743. PMID 23082175.
- [19] Chen J, Lariviere WR (2010). "The nociceptive and anti-nociceptive effects of bee venom injection and therapy: a double-edged sword". *Progress in Neurobiology*. **92** (2): 151–83. doi:10.1016/j.pneurobio.2010.06.006. PMC 2946189. PMID 20558236.
- [20] McGowan K, Kane A, Asarkof N, Wicks J, Guerina V, Kellum J, Baron S, Gintzler AR, Donowitz M (August 1983). "Entamoeba histolytica causes intestinal secretion: role of serotonin". *Science*. **221** (4612): 762–4. Bibcode:1983Sci...221..762M. doi:10.1126/science.6308760. PMID 6308760.
- [21] Feldman JM, Lee EM (October 1985). "Serotonin content of foods: effect on urinary excretion of 5-hydroxyindoleacetic acid". *The American Journal of Clinical Nutrition*. **42** (4): 639–43. doi:10.1093/ajcn/42.4.639. PMID 2413754.
- [22] Ramakrishna A, Giridhar P, Ravishankar GA (2011). "Phytoserotonin: a review". *Plant Signaling & Behavior*. **6** (6): 800–9. doi:10.4161/psb.6.6.15242. PMC 3218476. PMID 21617371.
- [23] Srinivasan S, Sadegh L, Elle IC, Christensen AG, Faergeman NJ, Ashrafi K (June 2008). "Serotonin regulates *C. elegans* fat and feeding through independent molecular mechanisms". *Cell Metabolism*. **7** (6): 533–44. doi:10.1016/j.cmet.2008.04.012. PMC 2495008. PMID 18522834.
- [24] Kravitz EA (September 1988). "Hormonal control of behavior: amines and the biasing of behavioral output in lobsters". *Science*. **241** (4874): 1775–81. Bibcode:1988Sci...241.1775K. doi:10.1126/science.2902685. PMID 2902685.
- [25] Beattie DT, Smith JA (May 2008). "Serotonin pharmacology in the gastrointestinal tract: a review". *Naunyn-Schmiedeberg's Archives of Pharmacology*. **377** (3): 181–203. doi:10.1007/s00210-008-0276-9. PMID 18398601. S2CID 32820765.
- [26] Sangkuhl K, Klein TE, Altman RB (November 2009). "Selective serotonin reuptake inhibitors pathway". *Pharmacogenetics and Genomics*. **19** (11): 907–9. doi:10.1097/FPC.0b013e32833132cb. PMC 2896866. PMID 19741567.
- [27] Hannon J, Hoyer D (December 2008). "Molecular biology of 5-HT receptors". *Behavioural Brain Research*. **195** (1): 198–213. doi:10.1016/j.bbr.2008.03.020. PMID 18571247. S2CID 46043982.
- [28] Zhou M, Engel K, Wang J (January 2007). "Evidence for significant contribution of a newly identified monoamine transporter (PMAT) to serotonin uptake in the human brain". *Biochemical Pharmacology*. **73** (1): 147–54. doi:10.1016/j.bcp.2006.09.008. PMC 1828907. PMID 17046718.
- [29] Paulmann N, Grohmann M, Voigt JP, Bert B, Vowinkel J, Bader M, Skelin M, Jevsek M, Fink H, Rupnik M, Walther DJ (October 2009). O'Rahilly S (ed.). "Intracellular serotonin modulates insulin secretion from pancreatic beta-cells by protein serenylation". *PLOS Biology*. **7** (10): e1000229. doi:10.1371/journal.pbio.1000229. PMC 2760755. PMID 19859528.
- [30] Walther DJ, Peter JU, Winter S, Höltje M, Paulmann N, Grohmann M, Vowinkel J, Alamo-Bethencourt V, Wilhelm CS, Ahnert-Hilger G, Bader M (December 2003). "Serenylation of small GTPases is a signal transduction pathway that triggers platelet alpha-granule release". *Cell*. **115** (7): 851–62. doi:10.1016/S0092-8674(03)01014-6. PMID 14697203. S2CID 16847296.
- [31] Watts SW, Priestley JR, Thompson JM (May 2009). "Serenylation of vascular proteins important to contraction". *PLOS ONE*. **4** (5): e5682. Bibcode:2009PLoS...4.5682W.

- doi:10.1371/journal.pone.0005682. PMC 2682564. PMID 19479059.*
- [32] Roth BL, Driscol J (12 January 2011). "PDSP Ki Database". Psychoactive Drug Screening Program (PDSP). University of North Carolina at Chapel Hill and the United States National Institute of Mental Health. Archived from the original on 8 November 2013. Retrieved 17 December 2013.
- [33] Bortolozzi A, Diaz-Mataix L, Scorza MC, Celada P, Artigas F (December 2005). "The activation of 5-HT receptors in prefrontal cortex enhances dopaminergic activity". *Journal of Neurochemistry*. **95** (6): 1597–607. *doi:10.1111/j.1471-4159.2005.03485.x. hdl:10261/33026. PMID 16277612.*
- [34] Moro C, Edwards L, Chess-Williams R (November 2016). "2Areceptor enhancement of contractile activity of the porcine urothelium and lamina propria". *International Journal of Urology*. **23** (11): 946–951. *doi:10.1111/iju.13172. PMID 27531585.*
- [35] "Von Economo neuron – NeuronBank". [neuronbank.org.](http://neuronbank.org/) [unreliable medical source?]
- [36] Millan MJ, Gobert A, Lejeune F, et al. (September 2003). "The novel melatonin agonist agomelatine (S20098) is an antagonist at 5-hydroxytryptamine2C receptors, blockade of which enhances the activity of frontocortical dopaminergic and adrenergic pathways". *The Journal of Pharmacology and Experimental Therapeutics*. **306** (3): 954–64. *doi:10.1124/jpet.103.051797. PMID 12750432. S2CID 18753440.*
- [37] Gonzalez R, Chávez-Pascacio K, Meneses A (September 2013). "Role of 5-HT5A receptors in the consolidation of memory". *Behavioural Brain Research*. **252**: 246–51. *doi:10.1016/j.bbr.2013.05.051. PMID 23735322. S2CID 140204585.*
- [38] Nautiyal KM, Hen R (2017). "Serotonin receptors in depression: from A to B". *F1000Research*. **6**: 123. *doi:10.12688/f1000research.9736.1. PMC 5302148. PMID 28232871.*
- [38] Frazer A, Hensler JG (1999). "Understanding the neuroanatomical organization of serotonergic cells in the brain provides insight into the functions of this neurotransmitter". In Siegel GJ, Agranoff, Bernard W, Fisher SK, Albers RW, Uhler MD (eds.). *Basic Neurochemistry (Sixth ed.)*. Lippincott Williams & Wilkins. ISBN 978-0-397-51820-3. In 1964, Dahlstrom and Fuxe (discussed in [2]), using the Falck-Hillarp technique of histofluorescence, observed that the majority of serotonergic soma are found in cell body groups, which previously had been designated as the Raphe nuclei.
- [39] Binder MD, Hirokawa N (2009). *encyclopedia of neuroscience*. Berlin: Springer. p. 705. ISBN 978-3-540-23735-8.
- [40] The raphe nuclei group of neurons are located along the brain stem from the labels 'Mid Brain' to 'Oblongata', centered on the pons. (See relevant image.)
- [38] Müller CP, Jacobs BL, eds. (2009). *Handbook of the behavioral neurobiology of serotonin (1st ed.)*. London: Academic. pp. 51–59. ISBN 9780123746344.
- [41] Sommer C (2009). "Serotonin in Pain and Pain Control". In Müller CP, Jacobs BL (eds.). *Handbook of the behavioral neurobiology of serotonin (1st ed.)*. London: Academic. pp. 457–460. ISBN 9780123746344.
- [42] Hensler JG (2009). "Serotonin in Mood and Emotions". In Müller CP, Jacobs BL (eds.). *Handbook of the behavioral neurobiology of serotonin (1st ed.)*. London: Academic. pp. 367–399. ISBN 9780123746344.
- [43] Andrews PW, Bharwani A, Lee KR, Fox M, Thomson JA (April 2015). "Is serotonin an upper or a downer? The evolution of the serotonergic system and its role in depression and the antidepressant response". *Neuroscience and Biobehavioral Reviews*. **51**: 164–88. *doi:10.1016/j.neubiorev.2015.01.018. PMID 25625874. S2CID 23980182.*
- [44] Stahl SM, Mignon L, Meyer JM (March 2009). "Which comes first: atypical antipsychotic treatment or cardiometabolic risk?". *Acta Psychiatrica Scandinavica*. **119** (3): 171–9. *doi:10.1111/j.1600-0447.2008.01334.x. PMID 19178394. S2CID 24035040.*
- [45] Buckland PR, Hoogendoorn B, Guy CA, Smith SK, Coleman SL, O'Donovan MC (March 2005). "Low gene expression conferred by association of an allele of the 5-HT2C receptor gene with antipsychotic-induced weight gain". *The American Journal of Psychiatry*. **162** (3): 613–5. *doi:10.1176/appi.ajp.162.3.613. PMID 15741483.*
- [46] Holmes MC, French KL, Seckl JR (June 1997). "Dysregulation of diurnal rhythms of serotonin 5-HT2C and corticosteroid receptor gene expression in the hippocampus with food restriction and glucocorticoids". *The Journal of Neuroscience*. **17** (11): 4056–65. *doi:10.1523/JNEUROSCI.17-11-04056.1997. PMC 6573558. PMID 9151722.*
- [47] Leibowitz SF (1990). "The role of serotonin in eating disorders". *Drugs. 39 Suppl 3*: 33–48. *doi:10.2165/00003495-199000393-00005. PMID 2197074. S2CID 8612545.*
- [48] McGuire, Michael (2013) "Believing, the neuroscience of fantasies, fears, and confictions" (Prometheus Books)
- [49] Caspi N, Modai I, Barak P, Waisbord A, Zbarsky H, Hirschmann S, Ritsner M (March 2001). "Pindolol augmentation in aggressive schizophrenic patients: a double-blind crossover randomized study". *International Clinical Psychopharmacology*. **16** (2): 111–5. *doi:10.1097/00004850-200103000-00006. PMID 11236069. S2CID 24822810.*
- [50] Ito Z, Aizawa I, Takeuchi M, Tabe M, Nakamura T (December 1975). "[Proceedings: Study of gastrointestinal motility using an extraluminal force transducer. 6. Observation of gastric and duodenal motility using synthetic motilin]". *Nihon Heikatsukin Gakkai Zasshi*. **11** (4): 244–6. *PMID 1232434.*
- [51] Lesch KP, Bengel D, Heils A, Sabol SZ, Greenberg BD, Petri S, Benjamin J, Müller CR, Hamer DH, Murphy DL (November 1996). "Association of anxiety-related traits with a polymorphism in the serotonin transporter gene regulatory region". *Science*. **274** (5292): 1527–31. *Bibcode:1996Sci...274.1527L.*

- doi:10.1126/science.274.5292.1527. PMID 8929413.*
S2CID 35503987.
- [52] Chilmonczyk Z, Bojarski AJ, Pilc A, Sylte I (August 2015). "Functional Selectivity and Antidepressant Activity of Serotonin 1A Receptor Ligands". *International Journal of Molecular Sciences*. **16** (8): 18474–506.
doi:10.3390/ijms160818474. PMC 4581256.
PMID 26262615.
- [53] Blier P, El Mansari M (2013). "Serotonin and beyond: therapeutics for major depression". *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*. **368** (1615): 20120536.
doi:10.1098/rstb.2012.0536. PMC 3638389.
PMID 23440470.
- [54] Eissa, Nermin; Al-Houqani, Mohammed; Sadeq, Adel; Ojha, Shreesh K.; Sasse, Astrid; Sadek, Bassem (16 May 2018). "Current Enlightenment About Etiology and Pharmacological Treatment of Autism Spectrum Disorder". *Frontiers in Neuroscience*. **12**: 304.
doi:10.3389/fnins.2018.00304. ISSN 1662-4548.
PMC 5964170. PMID 29867317.
- [55] Rang HP (2003). *Pharmacology*. Edinburgh: Churchill Livingstone. p. 187. ISBN 978-0-443-07145-4.
- [56] de Wit R, Aapro M, Blower PR (September 2005). "Is there a pharmacological basis for differences in 5-HT3-receptor antagonist efficacy in refractory patients?". *Cancer Chemotherapy and Pharmacology*. **56** (3): 231–8.
doi:10.1007/s00280-005-1033-0. PMID 15838653.
S2CID 27576150.
- [57] Frost M, Andersen TE, Yadav V, Brixen K, Karsenty G, Kassem M (March 2010). "Patients with high-bone-mass phenotype owing to Lrp5-T253I mutation have low plasma levels of serotonin". *Journal of Bone and Mineral Research*. **25** (3): 673–5. doi:10.1002/jbmr.44.
PMID 20200960.
- [58] Rosen CJ (February 2009). "Breaking into bone biology: serotonin's secrets". *Nature Medicine*. **15** (2): 145–6.
doi:10.1038/nm0209-145. PMID 19197289.
S2CID 5489589.
- [59] Mödder UI, Achenbach SJ, Amin S, Riggs BL, Melton LJ, Khosla S (February 2010). "Relation of serum serotonin levels to bone density and structural parameters in women". *Journal of Bone and Mineral Research*. **25** (2): 415–22. doi:10.1359/jbmr.090721. PMC 3153390.
PMID 19594297.
- [60] Frost M, Andersen T, Gossiel F, Hansen S, Bollerslev J, van Hul W, Eastell R, Kassem M, Brixen K (August 2011). "Levels of serotonin, sclerostin, bone turnover markers as well as bone density and microarchitecture in patients with high-bone-mass phenotype due to a mutation in Lrp5". *Journal of Bone and Mineral Research*. **26** (8): 1721–8.
doi:10.1002/jbmr.376. PMID 21351148.
- [61] Kode A, Mosialou I, Silva BC, Rached MT, Zhou B, Wang J, Townes TM, Hen R, DePinho RA, Guo XE, Kousteni S (October 2012). "FOXO1 orchestrates the bone-suppressing function of gut-derived serotonin". *The Journal of Clinical Investigation*. **122** (10): 3490–503.
doi:10.1172/JCI64906. PMC 3461930. PMID 22945629.
- [62] Yadav VK, Balaji S, Suresh PS, Liu XS, Lu X, Li Z, Guo XE, Mann JJ, Balapure AK, Gershon MD, Medhamurthy R, Vidal M, Karsenty G, Ducy P (March 2010). "Pharmacological inhibition of gut-derived serotonin synthesis is a potential bone anabolic treatment for osteoporosis". *Nature Medicine*. **16** (3): 308–12.
doi:10.1038/nm.2098. PMC 2836724. PMID 20139991.
- [63] Ozanne SE, Hales CN (January 2004). "Lifespan: catch-up growth and obesity in male mice". *Nature*. **427** (6973): 411–2. Bibcode:2004Natur.427..411O.
doi:10.1038/427411b. PMID 14749819. S2CID 40256021.
- [64] Lewis DS, Bertrand HA, McMahan CA, McGill HC, Carey KD, Masoro EJ (October 1986). "Preweaning food intake influences the adiposity of young adult baboons". *J. Clin. Invest.* **78** (4): 899–905. doi:10.1172/JCI112678.
PMC 423712. PMID 3760191.
- [65] Hahn P (July 1984). "Effect of litter size on plasma cholesterol and insulin and some liver and adipose tissue enzymes in adult rodents". *J. Nutr.* **114** (7): 1231–4.
doi:10.1093/jn/114.7.1231. PMID 6376732.
- [66] Popa D, Léna C, Alexandre C, Adrien J (April 2008). "Lasting syndrome of depression produced by reduction in serotonin uptake during postnatal development: evidence from sleep, stress, and behavior". *The Journal of Neuroscience*. **28** (14): 3546–54.
doi:10.1523/JNEUROSCI.4006-07.2008. PMC 6671102.